

## GENERAL RECOMMENDATIONS AFTER SURGERY

1. The tampon should be removed from the mouth after 30 minutes.
2. Cold compresses after the procedure apply alternately for 2-3 minutes as often as possible first two days after surgery. Avoid hot drinks and food.
3. After the procedure, we take food after the local anesthesia has subsided.
4. Do not drink alcohol or smoke cigarettes on the day of the procedure.
5. Carefully treat the post-treatment site until healing.
6. Painkiller, if necessary. . . . . times per day for . . . . . days.
7. Mouthwash 2-3 times a day with ELUDRIL starting 24 hours after procedure.
8. Antibiotic. . . . . x. . . . times per day (every . . . . . hours 1 tablet) for. . . . days.
9. Protective agent e. g. TRILAC 2 x per day 2 tablets.
10. AESCIN 3 x per day 2 tablets from the day of treatment (optional).
11. Teeth in the postoperative area can be brushed after. . . . . days.
12. Suture removal in. . . . . days.
13. If a surgical dressing is applied, it **should not** be brushed. After removing the dressing, treat the wound as above.

## GENERAL RECOMMENDATIONS AFTER SURGERY

1. The tampon should be removed from the mouth after 30 minutes.
2. Cold compresses after the procedure apply alternately for 2-3 minutes as often as possible first two days after surgery. Avoid hot drinks and food.
3. After the procedure, we take food after the local anesthesia has subsided.
4. Do not drink alcohol or smoke cigarettes on the day of the procedure.
5. Carefully treat the post-treatment site until healing.
6. Painkiller, if necessary. . . . . times per day for . . . . . days.
7. Mouthwash 2-3 times a day with ELUDRIL starting 24 hours after procedure.
8. Antibiotic. . . . . x. . . . times per day (every . . . . . hours 1 tablet) for. . . . days.
9. Protective agent e. g. TRILAC 2 x per day 2 tablets.
10. AESCIN 3 x per day 2 tablets from the day of treatment (optional).
11. Teeth in the postoperative area can be brushed after. . . . . days.
12. Suture removal in. . . . . days.
13. If a surgical dressing is applied, it **should not** be brushed. After removing the dressing, treat the wound as above.